

# Oxygen Tension Regulates Pancreatic $\beta$ -Cell Differentiation Through Hypoxia-Inducible Factor 1 $\alpha$

Mylène Heinis,<sup>1</sup> Marie-Thérèse Simon,<sup>1</sup> Karine Ilc,<sup>2</sup> Nathalie M. Mazure,<sup>2</sup> Jacques Pouyssegur,<sup>2</sup> Raphael Scharfmann,<sup>1</sup> and Bertrand Duvillie<sup>1</sup>

**OBJECTIVE**—Recent evidence indicates that low oxygen tension ( $pO_2$ ) or hypoxia controls the differentiation of several cell types during development. Variations of  $pO_2$  are mediated through the hypoxia-inducible factor (HIF), a crucial mediator of the adaptive response of cells to hypoxia. The aim of this study was to investigate the role of  $pO_2$  in  $\beta$ -cell differentiation.

**RESEARCH DESIGN AND METHODS**—We analyzed the capacity of  $\beta$ -cell differentiation in the rat embryonic pancreas using two in vitro assays. Pancreata were cultured either in collagen or on a filter at the air/liquid interface with various  $pO_2$ . An inhibitor of the prolyl hydroxylases, dimethylxaloylglycine (DMOG), was used to stabilize HIF1 $\alpha$  protein in normoxia.

**RESULTS**—When cultured in collagen, embryonic pancreatic cells were hypoxic and expressed HIF1 $\alpha$  and rare  $\beta$ -cells differentiated. In pancreata cultured on filter (normoxia), HIF1 $\alpha$  expression decreased and numerous  $\beta$ -cells developed. During pancreas development, HIF1 $\alpha$  levels were elevated at early stages and decreased with time. To determine the effect of  $pO_2$  on  $\beta$ -cell differentiation, pancreata were cultured in collagen at increasing concentrations of  $O_2$ . Such conditions repressed HIF1 $\alpha$  expression, fostered development of *Ngn3*-positive endocrine progenitors, and induced  $\beta$ -cell differentiation by  $O_2$  in a dose-dependent manner. By contrast, forced expression of HIF1 $\alpha$  in normoxia using DMOG repressed *Ngn3* expression and blocked  $\beta$ -cell development. Finally, hypoxia requires hairy and enhancer of split (HES)1 expression to repress  $\beta$ -cell differentiation.

**CONCLUSIONS**—These data demonstrate that  $\beta$ -cell differentiation is controlled by  $pO_2$  through HIF1 $\alpha$ . Modifying  $pO_2$  should now be tested in protocols aiming to differentiate  $\beta$ -cells from embryonic stem cells. *Diabetes* 59:662–669, 2010

**A**t low oxygen tension ( $pO_2$ ), cells undergo adaptive changes, including increased angiogenesis and erythropoiesis, and a switch to glycolytic metabolism (1). The cellular response to hypoxia is tightly controlled by the hypoxia-inducible factor (HIF) complex, an ( $\alpha/\beta$ ) heterodimer (2). HIF is regulated in an  $O_2$ -dependent manner by hydroxylation of one of the three HIF $\alpha$  subunits (HIF1 $\alpha$ , HIF2 $\alpha$ ,

and HIF3 $\alpha$ ) (3,4). HIF1 $\alpha$  was the original HIF isoform identified by affinity purification using oligonucleotides from the erythropoitin (EPO) locus, whereas HIF2 $\alpha$  and HIF3 $\alpha$  were identified by homology searches. These  $\alpha$ -subunits are expressed differently and independently in different time frames (5–7). In normoxia, hydroxylation of the proline HIF1 $\alpha$  residues by prolyl hydroxylases leads to recognition and polyubiquitinylation by the Von Hippel-Lindau (VHL) E3 ligase complex, leading to proteasomal degradation of HIF1 $\alpha$  (8,9). On the other hand, in the hypoxic state, prolyl hydroxylase activity is decreased and HIF1 $\alpha$  is stabilized. During embryogenesis, cell oxygenation levels vary widely. In general,  $pO_2$  is considered to occur at 2–9%  $O_2$  decreasing to even lower levels in tissues such as kidney medulla, bone marrow, or thymus (10). Recent investigations of the role of hypoxia and HIF1 $\alpha$  during development of several organs showed that hypoxia decreases the differentiation of neural precursor cells, myogenic cells, adipocytes, and endothelial cells in an HIF1 $\alpha$ -dependent manner (11–13).

The pancreas originates from the dorsal and ventral regions of the foregut endoderm directly behind the stomach. Signals derived from adjacent mesodermal structures, notochord, dorsal aorta, and cardiac mesoderm induce initiation of pancreas development (14–17). Next, mesenchyme condenses around the underlying committed endoderm; mesenchyme is also involved in the control of pancreas development (18). Embryonic pancreatic epithelium contains progenitor cells that express the transcription factor pancreatic and duodenal homeobox factor one (PDX1), whose deletion leads to pancreatic agenesis (19). Other transcription factors are also important for pancreatic development (20). Interestingly, the transcription factor *Ngn3* is transiently expressed in pancreatic endocrine progenitor cells during development, and its deletion leads to the absence of the four endocrine cell types ( $\alpha$ ,  $\beta$ ,  $\delta$ , and PP) that produce glucagon, insulin, somatostatin, and pancreatic polypeptide, respectively (21).

Advances in the understanding of signals that control pancreatic development have been made using in vivo and in vitro reconstituted models (15,22–24). In the present study, we found that when rat embryonic pancreata were cultured in collagen gels, cells were hypoxic, HIF1 $\alpha$  was stabilized, *Ngn3* expression was not induced, and consequently,  $\beta$ -cell differentiation was arrested. However, increasing  $pO_2$  led to HIF1 $\alpha$  degradation, upregulation of *Ngn3* expression, and  $\beta$ -cell differentiation. Similarly, when pancreata were cultured on filter at the air-medium interface, where tissues are not subject to hypoxia, HIF1 $\alpha$  was not expressed, *Ngn3* was induced, and  $\beta$ -cells developed. Under such conditions, reduction of  $pO_2$  stabilized HIF1 $\alpha$  levels and decreased both *Ngn3* expression and  $\beta$ -cell development. Moreover, HIF1 $\alpha$  stabilization using prolyl-hydroxylase inhibitors repressed  $\beta$ -cell develop-

From <sup>1</sup>INSERM U845, Research Center Growth and Signaling, Université Paris Descartes, Hôpital Necker, Paris, France; and the <sup>2</sup>Institute of Developmental Biology and Cancer Research, University of Nice, Nice, France.

Corresponding author: Bertrand Duvillie, bertrand.duvillie@inserm.fr. Received 17 June 2009 and accepted 2 December 2009. Published ahead of print at <http://diabetes.diabetesjournals.org> on 15 December 2009. DOI: 10.2337/db09-0891.

© 2010 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

ment. We also found that expression of *Hes1*, a transcription factor involved in pancreas development (25), was induced both by hypoxia and by forced expression of HIF1 $\alpha$  in normoxia. Moreover, inhibition of the *Hes1* expression by a  $\gamma$ -secretase inhibitor in hypoxic conditions restored normal development of *Ngn3*-expressing progenitor cells. Finally, we found that HIF1 $\alpha$  is temporally regulated during pancreas development in vivo. Based on these findings, we hypothesized that differences in pO<sub>2</sub> were responsible for such changes in  $\beta$ -cell development. Thus, we propose that pO<sub>2</sub> is a regulator of pancreatic endocrine cell development, acting through HIF1 $\alpha$ .

## RESEARCH DESIGN AND METHODS

Pregnant Wistar rats were purchased from the Janvier breeding center (CERJ). The animals had free access to food pellets and water. The first day postcoitus was taken as embryonic day 0.5 (E0.5). At 13.5 days' gestation, pregnant female rats were killed by asphyxiation with carbon dioxide or by a lethal injection of ketamin/xylazin according to the guidelines of the French Animal Care Committee.

**Hypoxia in vivo.** Pregnant Wistar rats at 13.5 days postcoitus were subjected to hypoxia in the anaerobic workstation at 25°C and in a humidified atmosphere (95%) supplied with the gas mixture allowing it to adjust the hypoxic environment by gradually decreasing the O<sub>2</sub> level from 21 to 8 and 92% N<sub>2</sub>. The rats were provided with food and water ad libitum. The animal study protocols were conducted according to institutional guidelines for animal use.

**Dissection of pancreatic rudiments and organ culture.** Embryos were harvested at E13.5. The dorsal pancreatic bud was dissected as described previously (23). For cultures at the air-medium interface, the dorsal pancreatic rudiments were laid on Millicell culture plate inserts (Millipore) (24). DMOG (Interchim) was used at concentrations of 1 and 2 mmol/l, and the  $\gamma$ -secretase inhibitor XXI (Calbiochem) was used at a concentration of 250 nmol/l. L-mimosine (Sigma) was used at a concentration of 400  $\mu$ mol/l. For collagen cultures, pancreatic explants were embedded in 500- $\mu$ l collagen gels as previously described (23). All cultures were maintained at 37°C in humidified 5% CO<sub>2</sub> in a hypoxia chamber at various levels of pO<sub>2</sub>.

**Detection of hypoxia.** To detect hypoxic areas, the dorsal pancreatic rudiments were cultured for either 24 h or 7 days in collagen and at the air-medium interface. Pimonidazole (200  $\mu$ mol/l; Hypoxyprobe, NPI) was added to culture medium during the last 2 h of the culture period. Pancreatic explants were analyzed with an anti-Mab1 fluorescent antibody (NPI) according to the manufacturer's instructions.

**Immunohistochemistry.** Tissues were fixed in 10% formalin and processed for immunohistochemistry as previously described (23,26). The following antibodies were used: mouse anti-insulin (1:2,000; Sigma-Aldrich), rabbit anti-PDX1 (1:1,000) (26), rabbit anti-carboxypeptidase A (1:600; Biogenesis), mouse anti-E-cadherin (1:100; BD Biosciences), rabbit anti-HES1 (1:1,000 [a gift from Dr. Tetsuo Sudo, Pharmaceutical Research Laboratories, Toray Industries, Teburo, Kamakura, Japan]), mouse anti-BrdU (1:2; Amersham), rabbit anti-NGN3 (1:1,000) (27), and rabbit anti-HIF1 $\alpha$  (9). The fluorescent secondary antibodies were fluorescein-isothiocyanate anti-rabbit and Texas Red anti-mouse antibodies (1:200; Jackson Immunoresearch) and Alexa-fluor anti-rabbit antibody (1:400; Biogenex). For NGN3, revelation was performed using the vectastain ABC kit (Vector). For HIF1 $\alpha$ , revelation was performed with an anti-rabbit Dakocytomation Envision+ system HRP-DAB (Dako). Photographs were taken using a fluorescence microscope (Leica, Leitz DMRB) and digitized using a Hamamatsu C5810 cooled 3CCD camera. No signals were observed when the primary antibodies were omitted.

**Transferase-mediated dUTP nick-end labeling experiments.** The transferase-mediated dUTP nick-end labeling (TUNEL) procedure was performed using an in situ cell death detection kit (Roche) according to the manufacturer's instructions. Subsequently, E-cadherin (1:100; BD Biosciences) immunostaining was used to visualize the epithelium.

**Quantification.** To quantify the absolute numbers of insulin-expressing cells, all sections of each pancreatic rudiment were digitized. On every image, the surface area of insulin staining was quantified using IPLab Eval (version 3.2.4; Scanalytics), and the stained areas were tallied as previously described (24). Three to five rudiments were analyzed per condition.

To measure proliferation of early progenitors expressing PDX1, we counted the frequency of BrdU<sup>+</sup> nuclei among 1,000 PDX1<sup>+</sup> cells. The percentage of undifferentiated PDX1-expressing cells was then calculated. Three rudiments were analyzed per condition. Statistical significance was determined using Student's *t* test.

**In situ hybridization.** Tissues were fixed at 4°C in 4% paraformaldehyde, cryoprotected in 15% sucrose at 4°C overnight, embedded in 15% sucrose and 7.5% gelatin, and frozen in isopentane. Cryosections (14  $\mu$ m thick) were prepared. An *Ngn3* probe (726 bp) was used and in situ hybridization performed as previously described (24). No signal was obtained using a sense riboprobe.

**Real-time PCR.** Total RNA was purified using the Rneasy microkit (Qiagen). The cDNA was generated using Superscript reagents (Invitrogen), and the real-time PCR was performed on a 7300 real-time PCR system (Applied Biosystem) with an SYBR Green PCR master mix. The oligonucleotide sequences for RT-PCR are available on request.


**Western blotting.** For Western blot analysis, cells were lysed in Laemmli buffer. Proteins (20  $\mu$ g) were resolved by SDS/PAGE and electrophoretically transferred onto a polyvinylidene fluoride membrane (Biorad). After blocking with milk, membranes were probed with rabbit anti-HIF1 $\alpha$  or anti-HIF2 $\alpha$  antibodies (Novus) and mouse anti-actin (Sigma-Aldrich). Immunoreactive bands were visualized with the SuperSignal system (Pierce).

## RESULTS

**Pancreatic  $\beta$ -cell differentiation depends on cell oxygenation.** We cultured E13.5 rat pancreata in collagen. While acinar cells expressing carboxypeptidase-A developed, NGN3 expression was poorly induced and  $\beta$ -cells did not develop. However, when we cultured them on a filter at the air-medium interface, NGN3 expression was induced and insulin-positive cells developed (Fig. 1A). When pimonidazole staining was applied, we found that collagen-cultured tissues were hypoxic from day 1 to day 7, which was not the case for pancreata cultured at the air-medium interface (Fig. 1B).

To investigate the role of O<sub>2</sub> in  $\beta$ -cell differentiation, we cultured pancreatic explants in collagen with increased pO<sub>2</sub>. When pancreata were cultured at 21% O<sub>2</sub>, very few  $\beta$ -cells were detected on day 7 (Fig. 2A). At 60 and 80% O<sub>2</sub>, the surface of insulin staining increased 25.1- and 81.2-fold, respectively, compared with cultures at 21% O<sub>2</sub> (Fig. 2A). Next, we examined development of different types of endocrine cells ( $\alpha$ ,  $\beta$ , and  $\delta$  cells) using quantitative real-time PCR (qPCR). When pancreatic buds were cultured at 80% O<sub>2</sub>, glucagon-, insulin-, and somatostatin-mRNA steady-state levels were increased 8.3-, 175.1-, and 8.9-fold, respectively, compared with levels induced at 21% O<sub>2</sub> (supplementary Fig. S1, which can be found in an online appendix at <http://diabetes.diabetesjournals.org/cgi/content/full/db09-0891/DC1>). Thus, pancreatic endocrine cell development is controlled by pO<sub>2</sub>. Interestingly, oxygen tension had no effect on acinar cell development (Fig. 2A).


To determine whether the increased  $\beta$ -cell development was due to higher cell survival or increased differentiation, we compared the rates of cell apoptosis at day 1 in pancreata cultured at 21 and 80% O<sub>2</sub>. Results of TUNEL staining detected very few apoptotic cells in the epithelia of pancreata treated in either culture (Fig. 2B). Some apoptotic cells were found in the mesenchyme; however, the numbers of apoptotic cells were similar for tissues in both cultures. Next, we used qPCR to compare *Ngn3* expression in pancreata cultured at 21 and 80% O<sub>2</sub> for 1, 3, and 5 days. At all time points, increased pO<sub>2</sub> induced *Ngn3* expression (Fig. 2C). Immunohistochemistry analysis confirmed these results (supplementary Fig. S2). Altogether, these data indicate that increased oxygen tension activates  $\beta$ -cell differentiation as opposed to enhancing cell survival. **HIF1 $\alpha$  signaling pathway is induced by hypoxia in the pancreas.** In various cell types, hypoxia stabilizes HIF1 $\alpha$  and activates the HIF transcription complex. In our study, we used Western blot analysis to determine whether HIF1 $\alpha$  was stabilized in pancreata cultured in collagen.



**FIG. 1.** Differential development of  $\beta$ -cells between pancreata cultured in collagen and at the air-medium interface. **A:** E13.5 pancreata were cultured in collagen or on a filter at the air-medium interface for 3 or 7 days. At day 3, NGN3 was detected by immunohistochemistry (in brown). At day 7, anti-insulin (INS) (in red) and anti-carboxypeptidase A (CPA) (in green) antibodies were used to detect  $\beta$ - and acinar cell development, respectively. **B:** Hypoxia was analyzed in pancreata cultured in collagen and at the air-medium interface. After 1 or 7 days, pimonidazole was added to the medium for the last 2 h of the culture period and hypoxia was analyzed using anti-Mab1 antibodies (in green). Bar: 50  $\mu$ m. (A high-quality digital representation of this figure is available in the online issue.)


After one day in collagen culture, HIF1 $\alpha$  levels at 21% O<sub>2</sub> increased dramatically compared with those cultured at 80% O<sub>2</sub> (Fig. 3A). To confirm these results, we used immunohistochemistry. Both HIF1 $\alpha$  and pimonidazole were detected at low but not high oxygen tension (Fig. 3B). We also analyzed the expression of HIF1 $\alpha$  targets. Using qPCR, we found that expression of the glycolytic enzymes *Pgk1* and *LdhA*, as well as expression of *Glut1*, *Vegf*, and *Redd1*, increased 9.1-, 8.4-, 11.3-, 4.1-, 4.7-fold, respectively, when cultured at 21% O<sub>2</sub> compared with 80% O<sub>2</sub> (Fig. 3C). Taken together, these results demonstrate that the HIF1 $\alpha$  signaling pathway is active in collagen culture at 21% O<sub>2</sub>, but not at 80% O<sub>2</sub>.

**Stabilization of HIF1 $\alpha$  at normoxia mimicks the effects of hypoxia on  $\beta$ -cell differentiation.** To determine whether HIF1 $\alpha$  was responsible for the arrest of  $\beta$ -cell differentiation under hypoxic conditions, we developed an approach to stabilize HIF1 $\alpha$  under normoxic conditions. To this end, we first showed that when pancreata were cultured on a filter at the air-medium interface, low pO<sub>2</sub> induced HIF1 $\alpha$  but not HIF2 $\alpha$  accumulation




**FIG. 2.** Induction of  $\beta$ -cell differentiation by increasing pO<sub>2</sub> in collagen. **A:** Pancreata were cultured for 7 days at 21, 60, and 80% O<sub>2</sub>.  $\beta$ - and acinar cells were detected by immunohistochemistry with antibodies directed against insulin (INS) (in red) and carboxypeptidase A (CPA) (in green). Next, the absolute surface area occupied by insulin-positive cells was quantified. **B:** Apoptotic cells (in green) were detected using a TUNEL reaction assay in pancreata cultured for 1 day at 21 or 80% O<sub>2</sub>. The epithelial cells were detected using an anti-E-cadherin antibody (in red). Note the absence of apoptotic cells within the epithelium. **C:** Real-time PCR quantification of *Ngn3* mRNA in pancreata cultured at 21% (□) or 80% (■) O<sub>2</sub> at different time points. Gene expression is presented as a percentage of the highest sample. Each point represents the mean  $\pm$  SEM of three individual data pools. \**P* < 0.05; \*\**P* < 0.01. (A high-quality digital representation of this figure is available in the online issue.)

(Fig. 4A and supplementary Fig. S3) and increased transcription of the HIF1 $\alpha$  target genes *Pgk1*, *LdhA*, *Glut1*, *Vegf*, and *Redd1* (supplementary Fig. S4). HIF1 $\alpha$  was expressed in the pimonidazole-positive areas (supplementary Fig. S5A), as was the case when pancreata were cultured in collagen (Fig. 3B). Moreover, under hypoxic conditions, *Ngn3* expression decreased both in mRNA




**FIG. 3.** The HIF1 $\alpha$  signaling pathway is induced by hypoxia in the pancreas. **A:** Protein extracts from pancreata cultured in collagen for 1 day at 21 or 80% O<sub>2</sub> were analyzed by Western blot to quantify HIF1 $\alpha$  expression and  $\beta$ -actin (loading control). **B:** Pancreata were cultured for 1 day at 21 or 80% O<sub>2</sub> and pimonidazole was added in the culture medium during the last 2 h of the culture. HIF1 $\alpha$ -positive cells were detected by immunohistochemistry (in brown). On serial sections, hypoxia was analyzed using anti-Mab1 antibodies (in green). Note the topographical relationship between HIF activation and pimonidazole adduct formation. Bar: 50  $\mu$ m. **C:** Expression of HIF1 $\alpha$  target genes *Pgk1*, *LdhA*, *Glut1*, *Vegf*, and *Redd1* was analyzed by qPCR in pancreata cultured in collagen for 1 day at 21% (□) or 80% (■) O<sub>2</sub>. Gene expression is presented as a percentage of the highest sample. Each point represents the mean  $\pm$  SEM of three individual pools. \* $P$  < 0.05; \*\* $P$  < 0.01. (A high-quality digital representation of this figure is available in the online issue.)

(Fig. 4B) and protein (supplementary Fig. S5B), and  $\beta$ -cell mass dramatically decreased (Fig. 4C). To evaluate the impact of oxygen tension on the proliferation of the precursor cells, we measured BrdU incorporation in pancreata cultured with 21 or 3% O<sub>2</sub> for 1 day (supplementary Fig. S6). There was a twofold decrease of BrdU incorporation in Pdx1-positive progenitors in the explants in hypoxia compared with normoxia. Thus, oxygen tension controls the proliferation of the progenitor cells. This effect should participate in the control of the number of NGN3-positive cells and the final  $\beta$ -cell mass. Next, we stabilized HIF1 $\alpha$  at normoxia with dimethylxaloylglycine (DMOG), a competitive inhibitor of prolyl hydroxylases (28). As shown in Fig. 4D, DMOG treatment caused HIF1 $\alpha$  accumulation and induction of the HIF1 $\alpha$  target genes in a dose-dependent manner (supplementary Fig. S7). On the contrary, DMOG did not stabilize HIF2 $\alpha$  (supplementary Fig. S3). Interestingly, under normoxic conditions, DMOG



**FIG. 4.** DMOG, an inhibitor of prolyl hydroxylases, stabilizes HIF1 $\alpha$  and represses  $\beta$ -cell differentiation. **A:** Pancreata at E13.5 were cultured at the air-medium interface for 1 day at 21 or 3% O<sub>2</sub>, and HIF1 $\alpha$  was analyzed by Western blot. **B:** Real-time PCR quantification of *Ngn3* mRNA in pancreata cultured for 1 day at 21% (■) or 3% (□) O<sub>2</sub>. Gene expression is presented as a percentage of the highest sample. Each point represents the mean  $\pm$  SEM of three individual pools. \* $P$  < 0.05. **C:** Pancreata were cultured for 7 days at the air-medium interface at 21 or 3% O<sub>2</sub>. Immunohistochemistry was used to detect  $\beta$ - and acinar cells using antibodies directed against insulin (in red) and carboxypeptidase A (in green). Next, the absolute surface occupied by insulin-positive cells was quantified. Black and gray bars represent the surface of insulin-positive cells in pancreata cultured at 21 and 3% O<sub>2</sub>, respectively. \*\*\* $P$  < 0.01. **D:** Pancreata at E13.5 were cultured at the air-medium interface for 1 day in the presence or absence of 1 or 2 mmol/l DMOG, and HIF1 $\alpha$  was analyzed by Western blot. **E:** Real-time PCR quantification of *Ngn3* mRNA in pancreata cultured for 1 day with 0 (■), 1 (□), or 2 (□) mmol/l DMOG at 21% O<sub>2</sub>. \*\* $P$  < 0.01. **F:** Pancreata cultured in the presence or absence of 1 mmol/l DMOG for 7 days were analyzed by immunohistochemistry. Acinar cells were detected using anti-carboxypeptidase A antibody (in green) and  $\beta$ -cells using anti-insulin antibody (in red). Next, the absolute surface occupied by insulin-positive cells was quantified. Gray and black bars represent the surface of insulin-positive cells in pancreata cultured at 21% O<sub>2</sub> with or without DMOG, respectively. \*\*\* $P$  < 0.001. (A high-quality digital representation of this figure is available in the online issue.)



**FIG. 5.** Regulation of *Hes1* expression by hypoxia. **A:** Real-time PCR quantification of *Hes1* mRNA in pancreata cultured at the air-medium interface with 21% (■) or 3% (□) O<sub>2</sub> for 1 day. \**P* < 0.05. **B:** Real-time PCR quantification of *Hes1* mRNA in pancreata cultured at the air-medium interface with 0 (■), 1 (□), or 2 (□) mmol/l DMOG for 1 day. \*\**P* < 0.01. **C:** Detection of *Ngn3* transcripts by in situ hybridization in pancreata cultured for 1 day at 21% O<sub>2</sub>, 21% O<sub>2</sub> with DMOG, or 3% O<sub>2</sub> with or without  $\gamma$ -secretase inhibitor treatment. Epithelium is circled in black in each panel. The absolute number of *Ngn3*-positive cells was quantified in each condition in the presence (■) or absence (□) of  $\gamma$ -secretase inhibitor treatment. \**P* < 0.05; \*\**P* < 0.01. (A high-quality digital representation of this figure is available in the online issue.)

treatment decreased *Ngn3* expression both in mRNA (Fig. 4E) and protein (supplementary Fig. S8). We further validated such results using L-mimosine, another inhibitor of prolyl hydroxylases (29–31). L-mimosine treatment resulted in HIF1 $\alpha$  stabilization and repression of *Ngn3* expression (supplementary Fig. S9). In addition, quantification of insulin staining indicated a sixfold decrease in  $\beta$ -cell development after 7 days of culture with DMOG compared with a culture without DMOG (Fig. 4F). Taken together, these results indicate that HIF1 $\alpha$  represses *Ngn3* expression and, consequently, negatively regulates  $\beta$ -cell differentiation.

**Hypoxia regulates *Hes1* expression via HIF1 $\alpha$ .** Based on our present discovery that collagen-cultured tissues are hypoxic, we hypothesized that hypoxia could control the expression of *Hes1*, a transcription factor involved in pancreas development (25). To prove this, we cultured E13.5 rat pancreata on filter with 3 and 21% O<sub>2</sub> and quantified expression of *Hes1*. We found that *Hes1* expression is activated by hypoxia (Fig. 5A). Results were similar


when DMOG was added to pancreata cultured at normoxia (Fig. 5B). To investigate whether hypoxia inhibits  $\beta$ -cell differentiation by regulating *Hes1* expression, we cultured pancreata under hypoxic conditions in the presence and absence of a  $\gamma$ -secretase inhibitor that inhibits the cleavage of Notch. In several cell types, *Hes1* has been described as a target of the Notch-activated receptor (11,32), but such a finding remains to be demonstrated in the pancreas (33,34). We thus first tested the effect of  $\gamma$ -secretase inhibitor XXI (Calbiochem) on HES1. Cultures of pancreatic explants were performed in the presence or absence of  $\gamma$ -secretase inhibitor, and HES1 protein expression was visualized using immunohistochemistry.  $\gamma$ -Secretase inhibitor abolished HES1 protein expression (supplementary Fig. S10). Interestingly, whereas in the absence of the  $\gamma$ -secretase inhibitor hypoxia decreased *Ngn3* expression, the presence of the  $\gamma$ -secretase inhibitor restored *Ngn3* expression (Fig. 5C). To further compare the effects of hypoxia and DMOG treatment, we cultured pancreata with DMOG in the presence or absence of  $\gamma$ -secretase inhibitor. As shown in Fig. 5C,  $\gamma$ -secretase inhibitor also abolishes the repression of *Ngn3* in DMOG-treated explants. Thus, these results demonstrate that both hypoxia and DMOG require expression of *Hes1* to repress *Ngn3*.

**HIF1 $\alpha$  protein expression decreases during pancreatic development.** Immunohistochemistry results showed that at E11.5 and E12.5, HIF1 $\alpha$  is present in both pancreatic epithelium and mesenchyme (Fig. 6A). Interestingly, HIF1 $\alpha$  levels are reduced at E16.5. By E20.5, HIF1 $\alpha$  expression decreased sharply and was rarely detected (Fig. 6A) and pancreatic islet cells stained negative for HIF1 $\alpha$  (Fig. 6A). Finally, HIF1 $\alpha$  is absent in the adult pancreas (Fig. 6A). Such a specific temporal pattern of expression was confirmed using Western blot analysis (supplementary Fig. S11).

**Hypoxia decreases  $\beta$ -cell development in vivo.** To investigate whether pO<sub>2</sub> regulates endocrine differentiation in vivo, we exposed pregnant rats at 13.5 days postcoitus to hypoxia at 8% O<sub>2</sub> or normoxia at 21% O<sub>2</sub> during 24 h as previously described (9). We next dissected the embryonic pancreases at E14.5. Expression of NGN3 was examined using immunohistochemistry, and expression of the HIF1 $\alpha$  target genes were quantified using qPCR. We found that NGN3 expression was dramatically reduced in hypoxia compared with that in normoxia while the HIF1 $\alpha$  target genes were induced (Fig. 6B). We thus conclude that experimental hypoxia downregulates endocrine differentiation in vivo as well as in vitro.

## DISCUSSION

Previously, we showed that in vitro, mature  $\beta$ -cells develop efficiently from rat embryonic pancreas grown at the air-medium interface, whereas  $\beta$ -cell development is poor when the same tissue is grown in collagen (23,24). Our working hypothesis was that hypoxia occurred in collagen gel, activating the Notch pathway and inhibiting endocrine differentiation (23,24,35). In our present study, we used pimonidazole staining to demonstrate that pancreata cultured in collagen were hypoxic compared with cultures grown at the air-medium interface. We also found that in vitro, the level of HIF1 $\alpha$ —a factor involved in regulation of many hypoxic responses (36)—increased in the pancreas as O<sub>2</sub> tension decreased and that it repressed pancreatic  $\beta$ -cell differentiation. At the difference to HIF1 $\alpha$ , the factor



**FIG. 6.** HIF1 $\alpha$  protein expression during pancreatic development in vivo. **A:** Immunohistochemical analysis of HIF1 $\alpha$  protein expression (in brown) in pancreata at E11.5, E12.5, E16.5, E20.5, and adult phases. In the *left panel*, epithelium is circled in black. In the *right panel*, a pancreatic islet is circled in black. Antibodies against PDX1 (in green) and insulin (INS) (in red) were used on serial sections to visualize progenitors and  $\beta$ -cells, respectively. **B:** NGN3 expression is repressed in the pancreas of embryos exposed to hypoxia. Pregnant rats were maintained with 21 or 8% O<sub>2</sub> for 24 h. Animal weight did not differ between the hypoxic and normoxic groups. The pancreata were harvested at E14.5 and analyzed by immunohistochemistry and qPCR. Detection of NGN3 (brown);  $^{***}P < 0.01$ . Expression of HIF1 $\alpha$  target genes, *Pgk1*, *LdhA*, *Glut-1*, *Vegf-A*, and *Redd1* in pancreata in hypoxia ( $\square$ ) or normoxia ( $\blacksquare$ ).  $^{*}P < 0.05$ ;  $^{***}P < 0.001$ . (A high-quality digital representation of this figure is available in the online issue.)

HIF2 $\alpha$  was not detected in the pancreas even in stimulated culture conditions (supplementary Fig. 3) or in VHL mutant mice (37). Data from literature indicate that HIF2 $\alpha$  is expressed during embryonic development mainly in endothelial cells (7,38,39). It is thus possible that variations of HIF2 $\alpha$  in the pancreatic endothelial cells may have influenced the development of endocrine cells in our experiments (17). However, our results indicating that hypoxia and DMOG do not stabilize HIF2 $\alpha$  do not favor this hypothesis.

Oxygen is known to control cell differentiation in different tissues. For example, hypoxia blocks myogenic and neuronal differentiation (11), and it arrests preadipocytes in an undifferentiated state (12,40). Thus, hypoxia seems to play a major role in maintaining a pool of undifferentiated stem/progenitor cells—a process important for their proper amplification. The fact that oxygen tension is low during early human and rodent development, when progenitor cell expansion is taking place (41,42), further supports this hypothesis. Recently, Fraker et al. (43) proposed that pO<sub>2</sub> plays a role in  $\beta$ -cell differentiation based on results with embryonic pancreata cultured on a silicone-perfluorocarbon membrane at different O<sub>2</sub> concentrations. However, the mechanism by which O<sub>2</sub> mediated its effects was not elucidated.

Recent data indicate that hypoxia, via HIF1 $\alpha$ , regulates limb bud differentiation. In this tissue, Provot et al. (44) proposed that condensation of the mesenchyme leads to

local hypoxia that stabilizes HIF1 $\alpha$  levels. Pancreatic anatomy during development suggests that pancreatic endocrine cell development could be controlled in the same way. Here we show that in vivo, HIF1 $\alpha$  levels are high at E11.5–E12.5 in the undifferentiated rat epithelium and in the mesenchyme, indicating that the pancreas is hypoxic at this early stage. This finding suggests that hypoxia could maintain the pool of progenitor cells in vivo. Moreover, when pregnant rats at 13.5 days postcoitus were exposed to hypoxia for 24 h, the developing embryonic pancreata failed to express NGN3 (Fig. 6B). This experimental finding supports the hypothesis that oxygen is a limiting parameter for  $\beta$ -cell differentiation in vivo. Moreover, cultures of pancreata with different physiological oxygen tensions showed that O<sub>2</sub> is required for  $\beta$ -cell development in vitro (Fig. 4 and data not shown). In fact, pO<sub>2</sub> influences both the proliferation of the progenitor cells (supplementary Fig. S6) and development of  $\beta$ -cells (Figs. 1 and 4). This finding should be considered when in vitro models are used to characterize the molecular mechanisms that control the development of the pancreas. Finally, we also found that increasing pO<sub>2</sub> to nonphysiological levels can enhance the development of  $\beta$ -cells in vitro (data not shown). This finding suggests that oxygen could be used as a new tool to generate  $\beta$ -cells in vitro. Recent studies have described the pancreatic phenotype of mice with different deletions of VHL or HIF1 $\alpha$  (37,45–47). Interestingly, Cantley et al. (37) used a pPDX1-cre

recombinase (CRE) to delete VHL during embryonic life. They found that 12 weeks after birth, HIF1 $\alpha$  was increased but  $\beta$ -cell mass was not modified. The difference between these results and ours could be due to the fact that  $\beta$ -cell mass was analyzed 12 weeks after birth, while VHL deletion had not occurred in all targeted cells (37). It is thus possible that compensation occurred during this developmental period. Moreover, Cantley et al. (37) used Pdx1-CRE recombinase to delete VHL in pancreatic progenitors. However, we found that at E11.5 and E12.5, HIF1 $\alpha$  was expressed in PDX1-positive pancreatic progenitor cells but also in surrounding mesenchymal cells. This observation was found both in vivo and in vitro (Fig. 3B; supplementary Figs. 5A and 6). We thus cannot exclude the possibility that HIF1 $\alpha$ , in addition to its role in progenitor cells, modifies epithelio-mesenchymal interactions that are known to be of major importance for pancreas development (24,48,49). One possibility could be that hypoxia modifies the signals from the mesenchyme, e.g., the ones mediated by FGF10 or Delta-like 1. Neither FGF10 nor delta-like 1 mRNA steady-state levels significantly differ in the explants cultured in hypoxia compared with those in normoxia (data not shown). Further genetic studies using conditional knockout of HIF1 $\alpha$  in mice or siRNA in cultured pancreatic explants will be necessary to analyze whether the effects of hypoxia on the epithelium are direct or involve the pancreatic mesenchyme.

In the present study, we demonstrate that the level of expression of *Ngn3*, which is a basic helix-loop-helix transcription factor expressed in islet cell progenitors and which is necessary to initiate the endocrine differentiation program (21,50), was inversely related to HIF1 $\alpha$  levels. Interestingly, *Ngn3* expression and pancreatic endocrine cell development are tightly regulated by *Hes1*. For example, deletion of *Hes1*, which is thought to be a target of the activated cleaved Notch, led to precocious endocrine cell differentiation, depletion of the progenitor cell pool, and a decrease in pancreatic cell mass (25). In the present study, we used a  $\gamma$ -secretase inhibitor to block the Notch pathway. While  $\gamma$ -secretase inhibitors have been used with success for this purpose (11,51,52), other substrates of the  $\gamma$ -secretase inhibitor have been listed, such as ErbB1 (epidermal growth factor receptor [EGFR]) and ErbB4, that are also involved in pancreas development (53–55) and we thus cannot exclude that  $\gamma$ -secretase inhibitor also acts on other pathways. Recent data indicate that  $\gamma$ -secretase inhibitor is able to repress HES1 expression within pancreatic explants (56). Our data confirmed this effect (supplementary Fig. S10). Importantly, in the presence of  $\gamma$ -secretase inhibitor, the effects of hypoxia and DMOG on *Ngn3* expression were alleviated. We thus concluded that HES1 was necessary for these effects, but the exact role of the Notch receptor should be clarified in further studies. Moreover, our data indicate that hypoxia activates *Hes1* through HIF1 $\alpha$ . Recently, a direct link was found between hypoxia and Notch signaling, and it is now proposed that these two pathways interact to inhibit differentiation of myogenic and neuronal progenitors during early stages of embryogenesis (11,57). Gustafsson et al. (11) reported that pharmacological inhibition of Notch signaling by a  $\gamma$ -secretase inhibitor alleviates inhibition of differentiation under hypoxic conditions. In addition, they reported a direct physical interaction between HIF1 $\alpha$  and the intracellular domain of the activated Notch receptor; HIF1 $\alpha$  was recruited to Notch-responsive promoters upon Notch acti-

vation under hypoxic conditions (11). These data are consistent with our findings and suggest that the mechanism by which HIF1 $\alpha$  activates the expression of *Hes1* under hypoxic conditions also applies to pancreatic endocrine cell development. The HIF1 $\alpha$ -*Hes1* pathway is thus a good candidate for an explanation of the effects of hypoxia.

Based on our observations, we speculate that hypoxia induces HIF1 $\alpha$  in early pancreatic progenitor cells, delaying their differentiation into endocrine cells and thus preserving a sufficient pool of precursors necessary for attaining an adequate final  $\beta$ -cell mass. These findings will help shed light on the molecular mechanisms responsible for  $\beta$ -cell differentiation. In addition, they should help define new protocols for deriving insulin-producing cells, thereby improving options for cell replacement therapy in type 1 diabetes.

#### ACKNOWLEDGMENTS

M.H. received support from the French Ministry for Research and Technology. This study received support from the National Institutes of Health Beta Cell Biology Consortium (DK 072495-02 [to R.S.]), the 6th European Union Framework Program  $\beta$ -Cell Therapy Integrated Project (to R.S.), and the Association Française des Diabétiques (to R.S.).

This study was also supported by an Alfediam/sanofi-aventis grant (to B.D.). No other potential conflicts of interest relevant to this article were reported.

#### REFERENCES

1. Semenza GL. Life with oxygen. *Science* 2007;318:62–64
2. Semenza GL. Hypoxia-inducible factor 1: master regulator of O<sub>2</sub> homeostasis. *Curr Opin Genet Dev* 1998;8:588–594
3. Ivan M, Kondo K, Yang H, Kim W, Valiando J, Ohh M, Salic A, Asara JM, Lane WS, Kaelin WG Jr. HIF1 $\alpha$  targeted for VHL-mediated destruction by proline hydroxylation: implications for O<sub>2</sub> sensing. *Science* 2001;292:464–468
4. Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, Kriegsheim A, Hübner HF, Mukherji M, Schofield CJ, Maxwell PH, Pugh CW, Ratcliffe PJ. Targeting of HIF-1 $\alpha$  to the von Hippel-Lindau ubiquitylation complex by O<sub>2</sub>-regulated prolyl hydroxylation. *Science* 2001;292:468–472
5. Holmquist-Mengelbier L, Fredlund E, Lofstedt T, Noguera R, Navarro S, Nilsson H, Pietras A, Vallon-Christersson J, Borg A, Gradin K, Poellinger L, Pahlman S. Recruitment of HIF-1 $\alpha$  and HIF-2 $\alpha$  to common target genes is differentially regulated in neuroblastoma: HIF-2 $\alpha$  promotes an aggressive phenotype. *Cancer Cell* 2006;10:413–423
6. Rosenberger C, Mandriota S, Jurgensen JS, Wiesener MS, Horstrup JH, Frei U, Ratcliffe PJ, Maxwell PH, Bachmann S, Eckardt KU. Expression of hypoxia-inducible factor-1 $\alpha$  and -2 $\alpha$  in hypoxic and ischemic rat kidneys. *J Am Soc Nephrol* 2002;13:1721–1732
7. Tian H, McKnight SL, Russell DW. Endothelial PAS domain protein 1 (EPAS1), a transcription factor selectively expressed in endothelial cells. *Genes Dev* 1997;11:72–82
8. Berra E, Ginouves A, Pouyssegur J. The hypoxia-inducible-factor hydroxylases bring fresh air into hypoxia signalling. *EMBO Rep* 2006;7:41–45
9. Ginouves A, Ilc K, Macias N, Pouyssegur J, Berra E. PHDs overactivation during chronic hypoxia “desensitizes” HIF1 $\alpha$  and protects cells from necrosis. *Proc Natl Acad Sci U S A* 2008;105:4745–4750
10. Simon MC, Keith B. The role of oxygen availability in embryonic development and stem cell function. *Nat Rev Mol Cell Biol* 2008;9:285–296
11. Gustafsson MV, Zheng X, Pereira T, Gradin K, Jin S, Lundkvist J, Ruas JL, Poellinger L, Lendahl U, Bondesson M. Hypoxia requires notch signaling to maintain the undifferentiated cell state. *Dev Cell* 2005;9:617–628
12. Lin Q, Lee YJ, Yun Z. Differentiation arrest by hypoxia. *J Biol Chem* 2006;281:30678–30683
13. Diez H, Fischer A, Winkler A, Hu CJ, Hatzopoulos AK, Breier G, Gessler M. Hypoxia-mediated activation of Dll4-Notch-Hey2 signaling in endothelial progenitor cells and adoption of arterial cell fate. *Exp Cell Res* 2007;313:1–9

14. Deutsch G, Jung J, Zheng M, Lora J, Zaret KS. A bipotential precursor population for pancreas and liver within the embryonic endoderm. *Development* 2001;128:871–881
15. Hebrok M, Kim SK, Melton DA. Notochord repression of endodermal Sonic hedgehog permits pancreas development. *Genes Dev* 1998;12:1705–1713
16. Kim SK, Hebrok M, Melton DA. Pancreas development in the chick embryo. *Cold Spring Harb Symp Quant Biol* 1997;62:377–383
17. Lammert E, Cleaver O, Melton D. Induction of pancreatic differentiation by signals from blood vessels. *Science* 2001;294:564–567
18. Scharfmann R. Control of early development of the pancreas in rodents and humans: implications of signals from the mesenchyme. *Diabetologia* 2000;43:1083–1092
19. Ahlgren U, Jonsson J, Edlund H. The morphogenesis of the pancreatic mesenchyme is uncoupled from that of the pancreatic epithelium in IPF1/PDX1-deficient mice. *Development* 1996;122:1409–1416
20. Collombat P, Hecksher-Sorensen J, Serup P, Mansouri A. Specifying pancreatic endocrine cell fates. *Mech Dev* 2006;123:501–512
21. Gradwohl G, Dierich A, LeMeur M, Guillemot F. Neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas. *Proc Natl Acad Sci U S A* 2000;97:1607–1611
22. Gittes GK, Galante PE, Hanahan D, Rutter WJ, Debase HT. Lineage-specific morphogenesis in the developing pancreas: role of mesenchymal factors. *Development* 1996;122:439–447
23. Miralles F, Czernichow P, Scharfmann R. Follistatin regulates the relative proportions of endocrine versus exocrine tissue during pancreatic development. *Development* 1998;125:1017–1024
24. Attali M, Stetsyuk V, Basmaciogullari A, Aiello V, Zanta-Boussif MA, Duvillie B, Scharfmann R. Control of  $\beta$ -cell differentiation by the pancreatic mesenchyme. *Diabetes* 2007;56:1248–1258
25. Jensen J, Pedersen EE, Galante P, Hald J, Heller RS, Ishibashi M, Kageyama R, Guillemot F, Serup P, Madsen OD. Control of endodermal endocrine development by Hes-1. *Nat Genet* 2000;24:36–44
26. Duvillie B, Attali M, Aiello V, Quemeneur E, Scharfmann R. Label-retaining cells in the rat pancreas: location and differentiation potential in vitro. *Diabetes* 2003;52:2035–2042
27. Guillemain G, Filhoulaud G, Da Silva-Xavier G, Rutter GA, Scharfmann R. Glucose is necessary for embryonic pancreatic endocrine cell differentiation. *J Biol Chem* 2007;282:15228–15237
28. Irwin R, LaPres JJ, Kinser S, McCabe LR. Prolyl-hydroxylase inhibition and HIF activation in osteoblasts promotes an adipocytic phenotype. *J Cell Biochem* 2007;100:762–772
29. Baader E, Tschank G, Baringhaus KH, Burghard H, Gunzler V. Inhibition of prolyl 4-hydroxylase by oxalyl amino acid derivatives in vitro, in isolated microsomes and in embryonic chicken tissues. *Biochem J* 1994;300:525–530
30. McCaffrey TA, Pomerantz KB, Sanborn TA, Spokojny AM, Du B, Park MH, Folk JE, Lamberg A, Kivirikko KI, Falcone DJ, et al. Specific inhibition of eIF-5A and collagen hydroxylation by a single agent: antiproliferative and fibrosuppressive effects on smooth muscle cells from human coronary arteries. *J Clin Invest* 1995;95:446–455
31. Warnecke C, Griethe W, Weidemann A, Jurgensen JS, Willam C, Bachmann S, Ivashchenko Y, Wagner I, Frei U, Wiesener M, Eckardt KU. Activation of the hypoxia-inducible factor-pathway and stimulation of angiogenesis by application of prolyl hydroxylase inhibitors. *Faseb J* 2003;17:1186–1188
32. Ohtsuka T, Ishibashi M, Gradwohl G, Nakanishi S, Guillemot F, Kageyama R. Hes1 and Hes5 as notch effectors in mammalian neuronal differentiation. *Embo J* 1999;18:2196–2207
33. Hald J, Hjorth JP, German MS, Madsen OD, Serup P, Jensen J. Activated Notch1 prevents differentiation of pancreatic acinar cells and attenuate endocrine development. *Dev Biol* 2003;260:426–437
34. Ahnfelt-Ronne J, Hald J, Bodker A, Yassin H, Serup P, Hecksher-Sorensen J. Preservation of proliferating pancreatic progenitor cells by Delta-Notch signaling in the embryonic chicken pancreas. *BMC Dev Biol* 2007;7:63
35. Duvillie B, Attali M, Bounacer A, Ravassard P, Basmaciogullari A, Scharfmann R. The mesenchyme controls the timing of pancreatic  $\beta$ -cell differentiation. *Diabetes* 2006;55:582–589
36. Brahimi-Horn MC, Pouyssegur J. Oxygen, a source of life and stress. *FEBS Lett* 2007;581:3582–3591
37. Cantley J, Selman C, Shukla D, Abramov AY, Forstreuter F, Esteban MA, Claret M, Lingard SJ, Clements M, Harten SK, Asare-Anane H, Batterham RL, Herrera PL, Persaud SJ, Duchon MR, Maxwell PH, Withers DJ. Deletion of the von Hippel-Lindau gene in pancreatic beta cells impairs glucose homeostasis in mice. *J Clin Invest* 2009;119:125–135
38. Flamme I, Frohlich T, von Reutern M, Kappel A, Damert A, Risau W. HRF, a putative basic helix-loop-helix-PAS-domain transcription factor is closely related to hypoxia-inducible factor-1 alpha and developmentally expressed in blood vessels. *Mech Dev* 1997;63:51–60
39. Ema M, Taya S, Yokotani N, Sogawa K, Matsuda Y, Fujii-Kuriyama Y. A novel bHLH-PAS factor with close sequence similarity to hypoxia-inducible factor 1alpha regulates the VEGF expression and is potentially involved in lung and vascular development. *Proc Natl Acad Sci U S A* 1997;94:4273–4278
40. Yun Z, Maecker HL, Johnson RS, Giaccia AJ. Inhibition of PPAR gamma 2 gene expression by the HIF-1-regulated gene DEC1/Stral3: a mechanism for regulation of adipogenesis by hypoxia. *Dev Cell* 2002;2:331–341
41. Yochim JM, Mitchell JA. Intrauterine oxygen tension in the rat during progestation: its possible relation to carbohydrate metabolism and the regulation of nidation. *Endocrinology* 1968;83:706–713
42. Rodesch F, Simon P, Donner C, Jauniaux E. Oxygen measurements in endometrial and trophoblastic tissues during early pregnancy. *Obstet Gynecol* 1992;80:283–285
43. Fraker CA, Alvarez S, Papadopoulos P, Giraldo J, Gu W, Ricordi C, Inverardi L, Dominguez-Bendala J. Enhanced oxygenation promotes beta-cell differentiation in vitro. *Stem Cells* 2007;25:3155–3164
44. Provot S, Zinyk D, Gunes Y, Kathri R, Le Q, Kronenberg HM, Johnson RS, Longaker MT, Giaccia AJ, Schipani E. Hif-1alpha regulates differentiation of limb bud mesenchyme and joint development. *J Cell Biol* 2007;177:451–464
45. Gunton JE, Kulkarni RN, Yim S, Okada T, Hawthorne WJ, Tseng YH, Roberson RS, Ricordi C, O'Connell PJ, Gonzalez FJ, Kahn CR. Loss of ARNT/HIF1beta mediates altered gene expression and pancreatic-islet dysfunction in human type 2 diabetes. *Cell* 2005;122:337–349
46. Puri S, Cano DA, Hebrok M. A role for von Hippel-Lindau protein in pancreatic  $\beta$ -cell function. *Diabetes* 2009;58:433–441
47. Zehetner J, Danzer C, Collins S, Eckhardt K, Gerber PA, Ballschiemier P, Galvanovskis J, Shimomura K, Ashcroft FM, Thorens B, Rorsman P, Krek W. PVHL is a regulator of glucose metabolism and insulin secretion in pancreatic beta cells. *Genes Dev* 2008;22:3135–3146
48. Bhushan A, Itoh N, Kato S, Thierry JP, Czernichow P, Bellusci S, Scharfmann R. Fgf10 is essential for maintaining the proliferative capacity of epithelial progenitor cells during early pancreatic organogenesis. *Development* 2001;128:5109–5117
49. Miralles F, Czernichow P, Ozaki K, Itoh N, Scharfmann R. Signaling through fibroblast growth factor receptor 2b plays a key role in the development of the exocrine pancreas. *Proc Natl Acad Sci U S A* 1999;96:6267–6272
50. Gu G, Dubauskaite J, Melton DA. Direct evidence for the pancreatic lineage: NGN3+ cells are islet progenitors and are distinct from duct progenitors. *Development* 2002;129:2447–2457
51. Dahlqvist C, Blokzijl A, Chapman G, Falk A, Dannaeus K, Ibanez CF, Lendahl U. Functional Notch signaling is required for BMP4-induced inhibition of myogenic differentiation. *Development* 2003;130:6089–6099
52. Cras-Meneur C, Li L, Kopan R, Permutt MA. Presenilins, Notch dose control the fate of pancreatic endocrine progenitors during a narrow developmental window. *Genes Dev* 2009;23:2088–2101
53. Miettinen PJ, Huotari M, Koivisto T, Ustinov J, Palgi J, Rasilainen S, Lehtonen E, Keski-Oja J, Otonkoski T. Impaired migration and delayed differentiation of pancreatic islet cells in mice lacking EGF-receptors. *Development* 2000;127:2617–2627
54. Kritzik MR, Krahl T, Good A, Gu D, Lai C, Fox H, Sarvetnick N. Expression of ErbB receptors during pancreatic islet development and regrowth. *J Endocrinol* 2000;165:67–77
55. Cras-Meneur C, Elghazi L, Czernichow P, Scharfmann R. Epidermal growth factor increases undifferentiated pancreatic embryonic cells in vitro: a balance between proliferation and differentiation. *Diabetes* 2001;50:1571–1579
56. Miralles F, Lamotte L, Couton D, Joshi RL. Interplay between FGF10 and Notch signalling is required for the self-renewal of pancreatic progenitors. *Int J Dev Biol* 2006;50:17–26
57. Cejudo-Martin P, Johnson RS. A new notch in the HIF belt: how hypoxia impacts differentiation. *Dev Cell* 2005;9:575–576